

Lecture 2. Analysis on normed linear space (I)

S is open if $\forall x \in S. \exists \epsilon > 0. \text{ s.t. } B(x, \epsilon) \subseteq S.$

S is closed if complement open.

definition

线性赋范空间

some examples.

$(0, 1)$ open

$[0, 1]$ closed

$(0, 1]$?

converge. $\{x_n\} \rightarrow x.$ or $\lim_{n \rightarrow \infty} x_n = x.$ if

$\lim_{n \rightarrow \infty} |x - x_n| = 0$ can be changed to any norm. proof?

in $\mathbb{R}.$ if $x_n \rightarrow x$ in one norm, then $x_n \rightarrow x$ in any norm

Theorem: S is closed iff \forall sequence $\{x_n\} \subseteq S.$ may not true for infinite dimension.

$$x_n \rightarrow x \Rightarrow x \in S.$$

$$f(A) = \begin{cases} k & [0, \frac{1}{k}] \\ 0 & \text{o.w.} \end{cases} \quad \begin{aligned} \|f\|_1 &\rightarrow 0 \\ \|f\|_2 &\rightarrow 1 \\ \|f\|_3 &\rightarrow \infty \end{aligned}$$

S is bounded if $\exists M < \infty. \text{ s.t. } \forall x \in S. |x| < M$

S is compact if S is closed and bounded.

Another definition: S is compact if any open covers has ^a finite subcover

continuous: a function f is continuous if at $x.$

$$\forall \epsilon > 0. \exists \delta > 0. \text{ s.t.}$$

$$\forall y \in X \cap B(x, \delta). |f(y) - f(x)| < \epsilon.$$

Extreme value theorem (Weierstrass).

If S is compact and $f: S \rightarrow \mathbb{R}$ continuous.


then f is bounded and exists extreme value.

sufficient but not necessary. see $\sin x / \cos x$.


Local minimum

x^* is a local minimum of f . if

$$\exists \varepsilon > 0. \text{ s.t. } \forall x \in X \cap B(x^*, \varepsilon), f(x) \geq f(x^*)$$



why does a function have min?
not



$$f(x): [0, 1] \rightarrow \mathbb{R}$$

$$f\left(\frac{p}{q}\right) = \frac{p}{q}$$

local information unbounded.

↑
continuous


if $f: \mathbb{R} \rightarrow \mathbb{R}$? \mathbb{R} is not compact.

if $f(\pm\infty) = \infty$. $\{x: f(x) < M\}$ compact.

Differential

$f(x): \mathbb{R} \rightarrow \mathbb{R}$.

$$f(x_0 + \delta) \approx k \cdot \delta + f(x_0)$$



Differential is a linear approximation of a function.

single value: $y = kx + b$.

general linear space?

linear $f(a \cdot u + v) = a f(u) + f(v)$.

affine 仿射 线性: 旋转 + 扭曲

仿射: 线性 + 平移

any linear operator is a matrix A .

affine operator: $Ax + b$.

$$\begin{cases} x \in \mathbb{R}^n \\ A \in \mathbb{R}^{m \times n} \\ b \in \mathbb{R}^m \end{cases}$$

$$f(x) \approx A(x - x_0) + f(x_0)$$

linear operator.

Def. differentiable: $f: \mathbb{R}^n \rightarrow \mathbb{R}^m$: if \exists matrix $A \in \mathbb{R}^{m \times n}$

$$\lim_{\substack{x \rightarrow x_0 \\ x \in X}} \frac{\|f(x) - (A(x - x_0) + f(x_0))\|}{\|x - x_0\|} = 0. \quad A \text{ is differential of } f.$$

denoted by $Df(x_0) = A$ or $f'(x_0) = A$.

Given $f(x) = (f_1(x), f_2(x), \dots, f_m(x))^T$

$Df(x_0)_{ij} = \frac{\partial f_i}{\partial x_j}(x_0)$. remark: $x_0 \in \mathbb{R}^n$ is a vector.

↑
called
Jacobian matrix

$$\begin{bmatrix} \frac{\partial f_1}{\partial x_1}(x_0) & \dots & \frac{\partial f_1}{\partial x_n}(x_0) \\ \vdots & & \vdots \\ \frac{\partial f_m}{\partial x_1}(x_0) & \dots & \frac{\partial f_m}{\partial x_n}(x_0) \end{bmatrix}$$

In particular, if $f: \mathbb{R}^n \rightarrow \mathbb{R}$. $Df \in \mathbb{R}^{1 \times n}$

gradient 梯度. $\nabla f(x) = (Df)^T = \left(\frac{\partial f}{\partial x_1}(x), \dots, \frac{\partial f}{\partial x_n}(x) \right)^T$

Example. $\begin{cases} f(x) = kx + b. & f'(x) = k \\ F(x) = Ax + b & F'(x) = A \end{cases}$

$\begin{cases} f(x) = ax^2 & f'(x) = 2a. \\ F(x) = x^T A x & F'(x) = ? x^T (A + A^T) \end{cases}$

$= 2A$ if A symmetric.

$F(x) = \sum_{i,j} A_{ij} x_i x_j$ $\frac{\partial F}{\partial x_k} = \sum_{i,j} A_{ij} \left(x_i \frac{\partial x_j}{\partial x_k} + x_j \frac{\partial x_i}{\partial x_k} \right)$

$f, g: \mathbb{R}^n \rightarrow \mathbb{R}^m$ $h \triangleq f^T g$
 $Dh(x) = f(x)^T Dg(x) + g(x)^T Df(x)$ $= \sum_i A_{ik} x_i + \sum_j A_{kj} x_j$.

Chain rule.

If $f: X \subseteq \mathbb{R}^n \rightarrow \mathbb{R}^m$ differentiable at $x_0 \in X$. $g: Y \subseteq \mathbb{R}^m \rightarrow \mathbb{R}^p$

then $h \triangleq g(f(x))$ differentiable at x_0 . differentiable at $y_0 = f(x_0)$

$Dh(x_0) = Dg(y_0) Df(x_0)$. Dg, Df matrix. order !!!

